

## FIFRA 2(ee) Recommendation



**ADDING SUPPRESSION OF ALTERNARIA, DOWNY MILDEWS, PESTALOTIOPSIS, AND POWDERY MILDEWS IN ALL LABELED CROPS  
CLARIFYING ACCEPTABILITY OF IN-FURROW, DRENCH OR TRANSPLANT STARTER SOLUTION USE OF HOWLER IN ALL LABELED CROPS  
EPA REG. No. 91197-3-92488**

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA. All applicable directions, restrictions, precautions, and Conditions of Sale or Warranty on the EPA registered label are to be followed. Refer to container label for additional instructions. Always read and follow label directions.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Always read this FIFRA 2(ee) bulletin and the entire label on the product container before using this product.

### **DIRECTIONS FOR USE**

#### **For Suppression of Alternaria, Downy Mildews, Pestalotiopsis, and Powdery Mildews**

For suppression of *Alternaria* (*Alternaria* spp), Downy Mildews (*Pseudoperonospora* spp, *Plasmopara* spp, *Sclerophthora* spp, *Phytophthora* spp, *Peronospora* spp, *Plasmopara* spp, and *Bremia* spp), Pestalotiopsis (*Pestalotiopsis* spp), Powdery Mildews (*Podosphaera* spp, *Erysiphe* spp, *Sphaerotheca* spp, *Leveillula* spp, *Microsphaera* spp, *Phyllactinia* spp, *Uncinula* spp, *Oidium* spp), in all labeled crops, apply Howler® Fungicide at 5 lb/acre. Howler® Fungicide can be applied by incorporation, drench, drip applications, in-furrow, foliar or chemigation.

#### **For All Labeled Crops**

For all labeled crops, Howler® Fungicide may be applied as an in-furrow spray, drench or transplant starter solution at an application rate of 5 – 15 lbs Howler® Fungicide/acre or 67 – 100 oz Howler® Fungicide/100 gal water.