

**FIFRA 2(ee)
Recommendation**



**ADDING FOLIAR APPLICATION TO POME FRUIT, CONTROL OF *MONILINIA*,
BOTRYOSPHAERIA, SHOT-HOLE, *FUSICLADIUM*, *DIAPORTHE*, AND
MYCOSPHAERELLA ON SELECTED CROPS AND ADDING SUPPRESSION OF FIRE
BLIGHT ON APPLE AND SCAB ON APPLE AND PEAR**

EPA REG. No. 91197-3-92488

**FOR USE AND DISTRIBUTION IN THE STATE OF TEXAS ONLY
THIS 2(ee) EXPIRES ON DECEMBER 31, 2024**

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA. All applicable directions, restrictions, precautions, and Conditions of Sale or Warranty on the EPA registered label are to be followed. Refer to the container label for additional instructions. Always read and follow label directions.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Always read this FIFRA 2(ee) bulletin and the entire label on the product container before using this product.

DIRECTIONS FOR USE

Add foliar application as an option for pome fruit

04/2021

For control of *Monilinia* (*Monilinia* spp.), *Botryosphaeria* (*Botryosphaeria* spp.), and Shot-Hole (*Wilsonomyces* spp.) on stone fruits and tree nuts

CROP	APPLICATION METHOD	APPLICATION RATE
Stone Fruits including Apricots, Cherries, Nectarines, Peaches, Plums, and Prunes	Foliar Spray	5 – 15 lb Howler®/acre
CROP	APPLICATION METHOD	APPLICATION RATE
Tree Nuts including Almonds, Beech Nuts, Brazil Nuts, Butternuts, Cashews, Chestnuts, Filberts, Hickory Nuts, Macadamia Nuts, Pecans, Pistachios, and Walnuts	Foliar Spray	5 – 15 lb Howler®/acre

For control of Scab (*Fusicladium* spp.) on Pecan

CROP	APPLICATION METHOD	APPLICATION RATE
Pecan	Foliar Spray	5 – 15 lb Howler®/acre

For control of Melanose (*Diaporthe* spp.) and Greasy spot (*Mycosphaerella* spp.) on citrus

CROP	APPLICATION METHOD	APPLICATION RATE
Citrus	Foliar Spray	5 – 15 lb Howler®/acre

For suppression of Fire Blight (*Erwinia* spp.) on apple

Apply Howler® at 5-15 lbs/acre in mixtures or rotations with other effective fungicides. Howler® can be applied as a foliar spray or via chemigation. Use of an adjuvant is recommended.

For suppression of Secondary Scab (*Venturia* spp.) on Apple and Pear

Apply Howler® at 5-15 lbs/acre in mixtures or rotations with other effective fungicides. Applications at petal fall through cover sprays. Howler can be applied as a foliar spray or via chemigation. Use of an adjuvant is recommended.

©2021 AgBiome Innovations, Inc. All Rights Reserved