

FIFRA 2(ee) Recommendation



**ADDING SUPPRESSION OF ALTERNARIA, DOWNY MILDEWS, PESTALOTIOPSIS, AND POWDERY MILDEWS IN ALL LABELED CROPS
CLARIFYING ACCEPTABILITY OF IN-FURROW, DRENCH OR TRANSPLANT STARTER SOLUTION USE OF HOWLER IN ALL LABELED CROPS**

EPA REG. No. 91197-3-92488

**FOR USE AND DISTRIBUTION IN THE STATE OF TEXAS ONLY
THIS 2(ee) EXPIRES ON DECEMBER 31, 2024**

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA. All applicable directions, restrictions, precautions, and Conditions of Sale or Warranty on the EPA registered label are to be followed. Refer to container label for additional instructions. Always read and follow label directions.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Always read this FIFRA 2(ee) Recommendation and the entire label on the product container before using this product.

DIRECTIONS FOR USE

For Suppression of Alternaria, Downy Mildews, Pestalotiopsis, and Powdery Mildews in All Labeled Crops

For suppression of Alternaria (*Alternaria* spp), Downy Mildews (*Pseudoperonospora* spp, *Plasmopara* spp, *Sclerophthora* spp, *Phytophthora* spp, *Peronospora* spp, *Plasmopara* spp, and *Bremia* spp), Pestalotiopsis (*Pestalotiopsis* spp), Powdery Mildews (*Podosphaera* spp, *Erysiphe* spp, *Sphaerotheca* spp, *Leveillula* spp, *Microsphaera* spp, *Phyllactinia* spp, *Uncinula* spp, *Oidium* spp), in all labeled crops, apply Howler® at 5 lb/acre. Howler® can be applied by incorporation, drench, in-furrow, foliar or chemigation as specified by crop on the container label.

For All Labeled Crops

For all labeled crops under greenhouse, nursery or field conditions, Howler® may be applied as an in-furrow spray, drench or transplant starter solution at an application rate of 5 – 15 lbs Howler®/acre or 67 – 100 oz Howler®/100 gal water.