

**FIFRA 2(ee)  
Recommendation**

**ADDING CONTROL OF ASCOCHYTA, BOTRYOSPHAERIA, MONILINIA, PHOMOPSIS, GUIGNARDIA,  
SPUR BLIGHT and GUMMY STEM BLIGHT.  
ADDING SUPPRESSION OF RUSTS OF SELECTED CROPS**

**EPA REG. No. 91197-3-92488**

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA. All applicable directions, restrictions, precautions, and Conditions of Sale or Warranty on the EPA registered label are to be followed. Refer to the container label for additional instructions. Always read and follow label directions.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Always read this FIFRA 2(ee) bulletin and the entire label on the product container before using this product.

Not for use in California, New York or Texas

**DIRECTIONS FOR USE**

**For control of *Monilinia* (*Monilinia* spp.) and *Botryosphaeria* (*Botryosphaeria* spp.) on berries and small fruits (e.g., blueberry (highbush and lowbush), blackberry, raspberry, cranberry)**

**For control of *Phomopsis* (*Phomopsis* spp.) on grapes, berries and small fruits (e.g., blueberry (highbush and lowbush), blackberry, raspberry, cranberry)**

**For control of Black rot (*Guignardia* spp.) on grapes**

**For control of Spur blight (*Didymella*) on raspberry**

**For control of Ascochyta blight (*Ascochyta* spp.) on beans, peas and dry pulses**

**For control of Gummy Stem Blight (*Stagonosporopsis* spp.) on cucurbit vegetables**

**For suppression of Rusts (*Puccinia* spp.) on small fruits**

apply 5-7.5 lbs/A (80-120 oz/100 gal water) of Howler as a foliar spray or via chemigation. Use of an adjuvant is recommended. Apply in rotations or mixtures with other effective fungicides.

© 2021 AgBiome Innovations, Inc. All rights reserved.