

**FIFRA 2(ee)  
Recommendation**



**ADDING CONTROL OF ASCOCHYTA, BOTRYOSPHERIA, MONILINIA, PHOMOPSIS, GUIGNARDIA,  
SPUR BLIGHT and GUMMY STEM BLIGHT.  
ADDING SUPPRESSION OF RUSTS OF SELECTED CROPS**

**EPA REG. No. 91197-3-92488**

**FOR USE AND DISTRIBUTION IN THE STATE OF TEXAS ONLY  
THIS 2(ee) EXPIRES ON DECEMBER 31, 2024**

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA. All applicable directions, restrictions, precautions, and Conditions of Sale or Warranty on the EPA registered label are to be followed. Refer to the container label for additional instructions. Always read and follow label directions.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Always read this FIFRA 2(ee) bulletin and the entire label on the product container before using this product.

**DIRECTIONS FOR USE**

**For control of *Monilinia* (*Monilinia* spp.) and *Botryosphaeria* (*Botryosphaeria* spp.) on berries and small fruits (e.g., blueberry (highbush and lowbush), blackberry, raspberry, cranberry).**

<b>CROP</b>	<b>APPLICATION METHOD</b>	<b>APPLICATION RATE</b>
<b><i>Berries and Small Fruits</i></b> including Blackberries, Blueberries, Currants, Elderberries, Gooseberries, Huckleberries, Loganberries, Raspberries, Strawberries	Foliar Spray	5 – 15 lb Howler®/acre

**For control of Phomopsis (*Phomopsis* spp.) on grapes, berries and small fruits (e.g., blueberry (highbush and lowbush), blackberry, raspberry, cranberry).**

CROP	APPLICATION METHOD	APPLICATION RATE
<b><i>Berries and Small Fruits</i></b> including Blackberries, Blueberries, Currants, Elderberries, Gooseberries, Huckleberries, Loganberries, Raspberries, Strawberries, and Grapes	Foliar Spray	5 – 15 lb Howler®/acre

**For control of Black rot (*Guignardia* spp.) on grapes.**

CROP	APPLICATION METHOD	APPLICATION RATE
Grapes	Foliar Spray	5 – 15 lb Howler®/acre

**For control of Spur blight (*Didymella*) on raspberry.**

CROP	APPLICATION METHOD	APPLICATION RATE
Raspberry	Foliar Spray	5 – 15 lb Howler®/acre

**For control of Ascochyta blight (*Ascochyta* spp.) on Beans, Peas, and Dry Pulses.**

CROP	APPLICATION METHOD	APPLICATION RATE
<b><i>Beans, Peas, and Dry Pulses</i></b>	Foliar or Directed Spray	5 – 15 lb Howler®/acre

**For control of Gummy Stem Blight (*Stagonosporopsis* spp.) on cucurbit vegetables.**

CROP	APPLICATION METHOD	APPLICATION RATE
<b><i>Cucurbit Vegetables</i></b> including Cucumbers, Cantaloupe, Melons, Gourds, Pumpkins, Winter Squash, Squash, and Zucchini	Foliar Spray	5 – 15 lb Howler®/acre
	Greenhouse Chemigation	67 – 100 oz Howler®/100 gal water

**For suppression of Rusts (*Puccinia* spp.) on small fruits.**

Apply 5-7.5 lbs/A (80-120 oz/100 gal water) of Howler as a foliar spray or via chemigation. Use of an adjuvant is recommended. Apply in rotations or mixtures with other effective fungicides.

© 2021 AgBiome Innovations, Inc. All rights reserved.