

April 21, 2021

New York State Department
of Environmental Conservation
Division of Materials Management
Pesticide Product Registration

Doc id: 573481

**FIFRA 2(ee)
Recommendation****ADDING FOLIAR APPLICATION TO POME FRUIT, CONTROL OF *MONILINIA* AND
SHOT-HOLE ON SELECTED CROPS AND ADDING SUPPRESSION OF FIRE BLIGHT
ON APPLE AND SCAB ON APPLE and PEAR****EPA REG. No. 91197-3-92488
FOR USE AND DISTRIBUTION IN THE STATE OF NEW YORK ONLY**

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA. All applicable directions, restrictions, precautions, and Conditions of Sale or Warranty on the EPA registered label are to be followed. Refer to the container label for additional instructions. Always read and follow label directions.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Always read this FIFRA 2(ee) bulletin and the entire label on the product container before using this product.

DIRECTIONS FOR USE**Add foliar application to pome fruit****For control of *Monilinia* (*Monilinia* spp.) and Shot-Hole (*Wilsonomyces* spp.) on
stone fruits and tree nuts**

Follow rates and timings in the Application Rate table on product label.

For suppression of Fire Blight (*Erwinia* spp.) on apple

Apply Howler® at 5-15 lbs/acre in mixtures or rotations with other effective fungicides. Howler® can be applied as a foliar spray or via chemigation. Use of an adjuvant is recommended.

For suppression of Secondary Scab (*Venturia* spp.) on Apple and Pear

Apply Howler® at 5-15 lbs/acre in mixtures or rotations with other effective fungicides. Applications should be made at petal fall through cover sprays. Howler can be applied as a foliar spray or via chemigation. Use of an adjuvant is recommended.

©2021 AgBiome Innovations, Inc. All Rights Reserved