

**FIFRA 2(ee)  
Recommendation**



**ADDING FOLIAR APPLICATION ON POME FRUIT (EXCLUDING PEARS AND QUINCE), CONTROL OF *MONILINIA*, *BOTRYOSPHAERIA*, SHOT-HOLE, *FUSICLADIUM*, *DIAPORTHE*, AND *MYCOSPHAERELLA* ON SELECTED CROPS AND ADDING SUPPRESSION OF FIRE BLIGHT ON APPLE AND SCAB ON APPLE**

**FOR USE AND DISTRIBUTION IN THE STATE OF CALIFORNIA ONLY**

**EPA REG. No. 91197-3-92488**

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA. All applicable directions, restrictions, precautions, and Conditions of Sale or Warranty on the EPA registered label are to be followed. Refer to the container label for additional instructions. Always read and follow label directions.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Always read this FIFRA 2(ee) bulletin and the entire label on the product container before using this product.

**DIRECTIONS FOR USE**

**Add foliar as an application method for pome fruit (excluding pears and quince)**

**For control of *Monilinia* (*Monilinia* spp.), *Botryosphaeria* (*Botryosphaeria* spp.), and Shot-Hole (*Wilsonomyces* spp.) on stone fruits (excluding plums and prunes) and tree nuts**

Follow rates and timings in the Application Rate table on product label

**For control of Scab (*Fusicladium* spp.) on pecan**

Follow rates and timings in the Application Rate table on product label

**For control of Melanose (*Diaporthe* spp.) and Greasy spot (*Mycosphaerella* spp.) on lemons and limes**

Follow rates and timings in the Application Rate table on product label

**For suppression of Fire Blight (*Erwinia* spp.) on apple**

Apply Howler® at 5-15 lbs/acre in mixtures or rotations with other effective fungicides. Howler® can be applied as a foliar spray or via chemigation. Use of an adjuvant is recommended.

**For suppression of Secondary Scab (*Venturia* spp.) on apple**

Apply Howler® at 5-15 lbs/acre in mixtures or rotations with other effective fungicides. Applications at petal fall through cover sprays. Howler can be applied as a foliar spray or via chemigation. Use of an adjuvant is recommended.

©2021 AgBiome Innovations, Inc. All Rights Reserved